

# Proton decay matrix elements from chirally symmetric lattice QCD

- ▷ Paul Cooney, The University of Edinburgh, RBC-UKQCD collaboration
- ▷ Workshop on Underground Detectors Investigating Grand Unification (UDiG) at Brookhaven National Laboratory



What to measure on the lattice

Simulation Details


Results

Non Perturbative Renormalization

Summary and Outlook

GUT Discrimination

$$p \rightarrow m + l$$



For a generic decay channel, the partial decay width is:

$$\Gamma(p \rightarrow m + \bar{l}) = \left[ \frac{m_p}{32\pi^2} \left( 1 - \left( \frac{m_m}{m_p} \right)^2 \right)^2 \right] \left| \sum_i C^i W_0^i(p \rightarrow m + \bar{l}) \right|^2$$

For a generic decay channel, the partial decay width is:

$$\Gamma(p \rightarrow m + \bar{l}) = \left[ \frac{m_p}{32\pi^2} \left( 1 - \left( \frac{m_m}{m_p} \right)^2 \right)^2 \right] \left| \sum_i C^i W_0^i(p \rightarrow m + \bar{l}) \right|^2$$

For a generic decay channel, the partial decay width is:

$$\Gamma(p \rightarrow m + \bar{l}) = \left[ \frac{m_p}{32\pi^2} \left( 1 - \left( \frac{m_m}{m_p} \right)^2 \right)^2 \right] \left| \sum_i C^i W_0^i(p \rightarrow m + \bar{l}) \right|^2$$

For a generic decay channel, the partial decay width is:

$$\Gamma(p \rightarrow m + \bar{l}) = \left[ \frac{m_p}{32\pi^2} \left( 1 - \left( \frac{m_m}{m_p} \right)^2 \right)^2 \right] \left| \sum_i C^i W_0^i(p \rightarrow m + \bar{l}) \right|^2$$

The form factors can be related to a matrix element

$$P_L [W_0^i(q^2) - i \not{q} W_q^i(q^2)] u(k, s) = \langle m | \mathcal{O}^i | N \rangle$$

For a generic decay channel, the partial decay width is:

$$\Gamma(p \rightarrow m + \bar{l}) = \left[ \frac{m_p}{32\pi^2} \left( 1 - \left( \frac{m_m}{m_p} \right)^2 \right)^2 \right] \left| \sum_i C^i W_0^i(p \rightarrow m + \bar{l}) \right|^2$$

The form factors can be related to a matrix element

$$P_L [\color{red}{W_0^i(q^2)} - i \not{q} W_q^i(q^2)] u(k, s) = \langle m | \mathcal{O}^i | N \rangle$$



For a generic decay channel, the partial decay width is:

$$\Gamma(p \rightarrow m + \bar{l}) = \left[ \frac{m_p}{32\pi^2} \left( 1 - \left( \frac{m_m}{m_p} \right)^2 \right)^2 \right] \left| \sum_i C^i W_0^i(p \rightarrow m + \bar{l}) \right|^2$$

The form factors can be related to a matrix element

$$P_L [W_0^i(q^2) - i \not{q} W_q^i(q^2)] u(k, s) = \langle m | \mathcal{O}^i | N \rangle$$

For a generic decay channel, the partial decay width is:

$$\Gamma(p \rightarrow m + \bar{l}) = \left[ \frac{m_p}{32\pi^2} \left( 1 - \left( \frac{m_m}{m_p} \right)^2 \right)^2 \right] \left| \sum_i C^i W_0^i(p \rightarrow m + \bar{l}) \right|^2$$

The form factors can be related to a matrix element

$$P_L W_0^i(q^2) u(k, s) = \langle m | \mathcal{O}^i | N \rangle$$

For a generic decay channel, the partial decay width is:

$$\Gamma(p \rightarrow m + \bar{l}) = \left[ \frac{m_p}{32\pi^2} \left( 1 - \left( \frac{m_m}{m_p} \right)^2 \right)^2 \right] \left| \sum_i C^i W_0^i(p \rightarrow m + \bar{l}) \right|^2$$

The form factors can be related to a matrix element

$$P_L W_0^i(q^2) u(k, s) = \langle m | \mathcal{O}^i | N \rangle$$

The operators  $\mathcal{O}^i$  are given by

$$\begin{aligned} \mathcal{O}^{RL} &= \epsilon^{abc} u^{a,T}(x, t) C P_R d^b(x, t) P_L u^c(x, t) \\ \mathcal{O}^{LL} &= \epsilon^{abc} u^{a,T}(x, t) C P_L d^b(x, t) P_L u^c(x, t) \end{aligned}$$

Define a general operator of the form

$$\mathcal{O}^{\Gamma_i \Gamma_j} = \epsilon^{abc} u^a(x, t) C \Gamma_i d^b(x, t) \Gamma_j u^c(x, t)$$

Define a general operator of the form

$$\mathcal{O}^{\Gamma_i \Gamma_j} = \epsilon^{abc} u^a(x, t) C \Gamma_i d^b(x, t) \Gamma_j u^c(x, t)$$

where  $\Gamma_i$  are matrices with two spin indices, labelled by,

$$\begin{aligned} S &= 1 & P &= \gamma_5 \\ V &= \gamma_\mu & A_\mu &= \gamma_\mu \gamma_5 \\ T &= \frac{1}{2} \{ \gamma_\mu, \gamma_\nu \} & \tilde{T} &= \gamma_5 \frac{1}{2} \{ \gamma_\mu, \gamma_\nu \} \\ R &= P_R = \frac{1}{2} (1 + \gamma_5) & L &= P_L = \frac{1}{2} (1 - \gamma_5) \end{aligned}$$

Define a general operator of the form

$$\mathcal{O}^{\Gamma_i \Gamma_j} = \epsilon^{abc} u^a(x, t) C \Gamma_i d^b(x, t) \Gamma_j u^c(x, t)$$

where  $\Gamma_i$  are matrices with two spin indices, labelled by,

$$\begin{aligned} S &= 1 & P &= \gamma_5 \\ V &= \gamma_\mu & A_\mu &= \gamma_\mu \gamma_5 \\ T &= \frac{1}{2} \{ \gamma_\mu, \gamma_\nu \} & \tilde{T} &= \gamma_5 \frac{1}{2} \{ \gamma_\mu, \gamma_\nu \} \\ R &= P_R = \frac{1}{2} (1 + \gamma_5) & L &= P_L = \frac{1}{2} (1 - \gamma_5) \end{aligned}$$

Operators with this structure are also used later in nucleon correlation functions and in the non-perturbative renormalization

We could measure the matrix elements  $\langle m | \mathcal{O}^i | N \rangle$  directly

- ▶ Known as the *direct* method
- ▶ Three-point functions are required
- ▶ Computationally expensive

Alternatively can relate the three-point functions to two-point functions using Chiral Perturbation Theory

- ▶ Known as the *indirect* method
- ▶ Computationally cheaper
- ▶ Introduces an additional source of error

We could measure the matrix elements  $\langle m | \mathcal{O}^i | N \rangle$  directly

- ▶ Known as the *direct* method
- ▶ Three-point functions are required
- ▶ Computationally expensive

Alternatively can relate the three-point functions to two-point functions using Chiral Perturbation Theory

- ▶ Known as the *indirect* method
- ▶ Computationally cheaper
- ▶ Introduces an additional source of error



We could measure the matrix elements  $\langle m | \mathcal{O}^i | N \rangle$  directly

- ▶ Known as the *direct* method
- ▶ Three-point functions are required
- ▶ Computationally expensive

Alternatively can relate the three-point functions to two-point functions using Chiral Perturbation Theory

- ▶ Known as the *indirect* method
- ▶ Computationally cheaper
- ▶ Introduces an additional source of error

We could measure the matrix elements  $\langle m | \mathcal{O}^i | N \rangle$  directly

- ▶ Known as the *direct* method
- ▶ Three-point functions are required
- ▶ Computationally expensive

Alternatively can relate the three-point functions to two-point functions using Chiral Perturbation Theory

- ▶ Known as the *indirect* method
- ▶ Computationally cheaper
- ▶ Introduces an additional source of error

For  $p \rightarrow \pi^0 + e^+$ , the chiral perturbation theory gives

$$W_0^{RL}(p \rightarrow \pi^0 + e^+) = \alpha(1 + D + F)/\sqrt{2}f + \mathcal{O}(m_l^2/m_N^2)$$

$$W_0^{LL}(p \rightarrow \pi^0 + e^+) = \beta(1 + D + F)/\sqrt{2}f + \mathcal{O}(m_l^2/m_N^2)$$

For  $p \rightarrow \pi^0 + e^+$ , the chiral perturbation theory gives

$$W_0^{RL}(p \rightarrow \pi^0 + e^+) = \alpha(1 + D + F)/\sqrt{2}f + \mathcal{O}(m_l^2/m_N^2)$$

$$W_0^{LL}(p \rightarrow \pi^0 + e^+) = \beta(1 + D + F)/\sqrt{2}f + \mathcal{O}(m_l^2/m_N^2)$$

$\alpha$  and  $\beta$  are low energy constants from the chiral lagrangian

For  $p \rightarrow \pi^0 + e^+$ , the chiral perturbation theory gives

$$W_0^{RL}(p \rightarrow \pi^0 + e^+) = \alpha(1 + D + F)/\sqrt{2}f + \mathcal{O}(m_l^2/m_N^2)$$

$$W_0^{LL}(p \rightarrow \pi^0 + e^+) = \beta(1 + D + F)/\sqrt{2}f + \mathcal{O}(m_l^2/m_N^2)$$

$\alpha$  and  $\beta$  are low energy constants from the chiral lagrangian  
They can be calculated from two-point functions

$$\langle 0 | \mathcal{O}^{RL} | N \rangle = \alpha P_L u(k, s)$$

$$\langle 0 | \mathcal{O}^{LL} | N \rangle = \beta P_L u(k, s)$$

Define a class of two-point functions

$$f_{\Gamma_1\Gamma_2,\Gamma_3\Gamma_4}(t) = \sum_x \text{tr} \left[ \langle \mathcal{O}^{\Gamma_1\Gamma_2} \bar{\mathcal{O}}^{\Gamma_3\Gamma_4} \rangle P \right]$$

Define a class of two-point functions

$$f_{\Gamma_1 \Gamma_2, \Gamma_3 \Gamma_4}(t) = \sum_x \text{tr} \left[ \langle \mathcal{O}^{\Gamma_1 \Gamma_2} \bar{\mathcal{O}}^{\Gamma_3 \Gamma_4} \rangle P \right]$$

$$\mathcal{O}^{\Gamma_1 \Gamma_2} = \epsilon^{abc} u^a(x, t) C \Gamma_i d^b(x, t) \Gamma_j u^c(x, t)$$

Define a class of two-point functions

$$f_{\Gamma_1\Gamma_2,\Gamma_3\Gamma_4}(t) = \sum_x \text{tr} \left[ \langle \mathcal{O}^{\Gamma_1\Gamma_2} \bar{\mathcal{O}}^{\Gamma_3\Gamma_4} \rangle \textcolor{red}{P} \right]$$

$$\mathcal{O}^{\Gamma_1\Gamma_2} = \epsilon^{abc} u^a(x, t) C \Gamma_i d^b(x, t) \Gamma_j u^c(x, t)$$

$$\textcolor{red}{P} = \frac{1}{2}(1 + \gamma_4)$$



Define a class of two-point functions

$$f_{\Gamma_1\Gamma_2,\Gamma_3\Gamma_4}(t) = \sum_x \text{tr} \left[ \langle \mathcal{O}^{\Gamma_1\Gamma_2} \bar{\mathcal{O}}^{\Gamma_3\Gamma_4} \rangle P \right]$$

$$\mathcal{O}^{\Gamma_1\Gamma_2} = \epsilon^{abc} u^a(x, t) C \Gamma_i d^b(x, t) \Gamma_j u^c(x, t)$$

$$P = \frac{1}{2}(1 + \gamma_4)$$

Example: the proton correlation function

$$\sum_x \langle J_p(x, t) \bar{J}_p(0) \rangle = f_{PS,PS}(t)$$

Strategy:

- First find  $m_N$  from a correlated fit to the effective mass

$$m_{\text{eff}}(t) = \log \left( \frac{f_{PS,PS}(t)}{f_{PS,PS}(t+1)} \right) \rightarrow m_N \quad t \gg 0$$

- Then find  $G_N$  from a correlated fit to an effective amplitude

$$G_{N,\text{eff}} = \sqrt{2f_{PS,PS} e^{m_N t}} \rightarrow G_N \quad t \gg 0$$

- Finally to calculate  $\alpha$  and  $\beta$  we use a ratio of two-point functions

$$R_\alpha(t) = 2G_N \frac{f_{RL,PS}(t)}{f_{PS,PS}(t)} \rightarrow \alpha \quad R_\beta(t) = 2G_N \frac{f_{LL,PS}(t)}{f_{PS,PS}(t)} \rightarrow \beta$$

Strategy:

- First find  $m_N$  from a correlated fit to the effective mass

$$m_{\text{eff}}(t) = \log \left( \frac{f_{PS,PS}(t)}{f_{PS,PS}(t+1)} \right) \rightarrow m_N \quad t \gg 0$$

- Then find  $G_N$  from a correlated fit to an effective amplitude

$$G_{N,\text{eff}} = \sqrt{2f_{PS,PS} e^{m_N t}} \rightarrow G_N \quad t \gg 0$$

- Finally to calculate  $\alpha$  and  $\beta$  we use a ratio of two-point functions

$$R_\alpha(t) = 2G_N \frac{f_{RL,PS}(t)}{f_{PS,PS}(t)} \rightarrow \alpha \quad R_\beta(t) = 2G_N \frac{f_{LL,PS}(t)}{f_{PS,PS}(t)} \rightarrow \beta$$

Strategy:

- First find  $m_N$  from a correlated fit to the effective mass

$$m_{\text{eff}}(t) = \log \left( \frac{f_{PS,PS}(t)}{f_{PS,PS}(t+1)} \right) \rightarrow m_N \quad t \gg 0$$

- Then find  $G_N$  from a correlated fit to an effective amplitude

$$G_{N,\text{eff}} = \sqrt{2f_{PS,PS} e^{m_N t}} \rightarrow G_N \quad t \gg 0$$

- Finally to calculate  $\alpha$  and  $\beta$  we use a ratio of two-point functions

$$R_\alpha(t) = 2G_N \frac{f_{RL,PS}(t)}{f_{PS,PS}(t)} \rightarrow \alpha \quad R_\beta(t) = 2G_N \frac{f_{LL,PS}(t)}{f_{PS,PS}(t)} \rightarrow \beta$$

- ▶ Calculation is carried out on 2+1 flavour Domain Wall Fermion ensembles
  - ▶ Nearly exact chiral symmetry
  - ▶ Inverse lattice spacing  $a^{-1} = 1.73(3)$  GeV
- ▶ Two different lattice volumes
  - ▶  $V = 16^3 \times 32 \approx 1.8\text{fm}^3$
  - ▶  $V = 24^3 \times 64 \approx 2.7\text{fm}^3$
- ▶ One strange quark with mass  
 $am_s = 0.04$
- ▶ Two degenerate light quarks with masses  
 $am_{u/d} = 0.005^*, 0.01, 0.02$  or  $0.03$

- ▶ Calculation is carried out on 2+1 flavour Domain Wall Fermion ensembles
  - ▶ Nearly exact chiral symmetry
  - ▶ Inverse lattice spacing  $a^{-1} = 1.73(3)$  GeV
- ▶ Two different lattice volumes
  - ▶  $V = 16^3 \times 32 \approx 1.8\text{fm}^3$
  - ▶  $V = 24^3 \times 64 \approx 2.7\text{fm}^3$
- ▶ One strange quark with mass  
 $am_s = 0.04$
- ▶ Two degenerate light quarks with masses  
 $am_{u/d} = 0.005^*, 0.01, 0.02$  or  $0.03$

- ▶ Calculation is carried out on 2+1 flavour Domain Wall Fermion ensembles
  - ▶ Nearly exact chiral symmetry
  - ▶ Inverse lattice spacing  $a^{-1} = 1.73(3)$  GeV
- ▶ Two different lattice volumes
  - ▶  $V = 16^3 \times 32 \approx 1.8\text{fm}^3$
  - ▶  $V = 24^3 \times 64 \approx 2.7\text{fm}^3$
- ▶ One strange quark with mass  
 $am_s = 0.04$
- ▶ Two degenerate light quarks with masses  
 $am_{u/d} = 0.005^*, 0.01, 0.02$  or  $0.03$

- ▶ Calculation is carried out on 2+1 flavour Domain Wall Fermion ensembles
  - ▶ Nearly exact chiral symmetry
  - ▶ Inverse lattice spacing  $a^{-1} = 1.73(3)$  GeV
- ▶ Two different lattice volumes
  - ▶  $V = 16^3 \times 32 \approx 1.8\text{fm}^3$
  - ▶  $V = 24^3 \times 64 \approx 2.7\text{fm}^3$
- ▶ One strange quark with mass  
 $am_s = 0.04$
- ▶ Two degenerate light quarks with masses  
 $am_{u/d} = 0.005^*, 0.01, 0.02$  or  $0.03$



## Fitting

Fit by minimising a correlated  $\chi^2$

$$\chi^2(p) = \sum_{t,t'} [p_{\text{eff}}(t) - p] C_{tt'}^{-1} [p_{\text{eff}}(t') - p]$$

With correlation Matrix

$$C_{tt'} = \frac{1}{N_{\text{boot}}} \sum_{n=1}^{N_{\text{boot}}} \left[ p_{\text{eff}}^{(n)}(t) - \bar{p}_{\text{eff}}(t) \right] \left[ p_{\text{eff}}^{(n)}(t') - \bar{p}_{\text{eff}}(t') \right].$$

Bootstrap to get central value and errors

## Fitting

Fit by minimising a correlated  $\chi^2$

$$\chi^2(p) = \sum_{t,t'} [p_{\text{eff}}(t) - p] \mathbf{C}_{tt'}^{-1} [p_{\text{eff}}(t') - p]$$

With correlation Matrix

$$\mathbf{C}_{tt'} = \frac{1}{N_{\text{boot}}} \sum_{n=1}^{N_{\text{boot}}} \left[ p_{\text{eff}}^{(n)}(t) - \bar{p}_{\text{eff}}(t) \right] \left[ p_{\text{eff}}^{(n)}(t') - \bar{p}_{\text{eff}}(t') \right].$$

Bootstrap to get central value and errors

## Fitting

Fit by minimising a correlated  $\chi^2$



$$\chi^2(p) = \sum_{t,t'} [p_{\text{eff}}(t) - p] \mathbf{C}_{tt'}^{-1} [p_{\text{eff}}(t') - p]$$

With correlation Matrix



$$\mathbf{C}_{tt'} = \frac{1}{N_{\text{boot}}} \sum_{n=1}^{N_{\text{boot}}} \left[ p_{\text{eff}}^{(n)}(t) - \bar{p}_{\text{eff}}(t) \right] \left[ p_{\text{eff}}^{(n)}(t') - \bar{p}_{\text{eff}}(t') \right].$$

Bootstrap to get central value and errors



# Nucleon Mass and amplitude





# Nucleon Extrapolations



## Low energy constant: $\alpha$



Low energy constant:  $\beta$



## Systematic Errors

- ▶ **Finite volume errors**
- ▶ Chiral Extrapolation errors
- ▶ (Continuum Extrapolation errors)
- ▶ Errors in renormalisation
- ▶ (Error in Chiral Perturbation Theory)



## Systematic Errors

- ▶ Finite volume errors
- ▶ Chiral Extrapolation errors
- ▶ (Continuum Extrapolation errors)
- ▶ Errors in renormalisation
- ▶ (Error in Chiral Perturbation Theory)

## Systematic Errors

- ▶ Finite volume errors
- ▶ Chiral Extrapolation errors
- ▶ (Continuum Extrapolation errors)
- ▶ Errors in renormalisation
- ▶ (Error in Chiral Perturbation Theory)


## Systematic Errors

- ▶ Finite volume errors
- ▶ Chiral Extrapolation errors
- ▶ (Continuum Extrapolation errors)
- ▶ Errors in renormalisation
- ▶ (Error in Chiral Perturbation Theory)

## Systematic Errors


- ▶ Finite volume errors
- ▶ Chiral Extrapolation errors
- ▶ (Continuum Extrapolation errors)
- ▶ Errors in renormalisation
- ▶ (Error in Chiral Perturbation Theory)

## Finite Volume Error




► No noticeable effect

# Extrapolation Error



18%



17%

## NPR

- ▶ Non-perturbative MOM scheme renormalisation of the Rome-Southampton group
- ▶ The renormalised operators are

$$\mathcal{O}_{\text{ren}}^A = Z^{AB} \mathcal{O}_{\text{latt}}^B$$

- ▶ A and B label the spin structure, eg  $LL$
- ▶  $Z^{AB}$  is the mixing matrix
- ▶  $\mathcal{O}^{LL}$  and  $\mathcal{O}^{RL}$  mix with a 3rd operator  $\mathcal{O}^{A(LV)}$   
 $\Rightarrow Z^{AB}$  is a  $3 \times 3$  matrix
- ▶ Exponentially accurate chiral symmetry from Domain Wall Fermions should suppress operator mixing

## NPR

- ▶ Non-perturbative MOM scheme renormalisation of the Rome-Southampton group
- ▶ The renormalised operators are

$$\mathcal{O}_{\text{ren}}^A = Z^{AB} \mathcal{O}_{\text{latt}}^B$$

- ▶ A and B label the spin structure, eg  $LL$
- ▶  $Z^{AB}$  is the mixing matrix
- ▶  $\mathcal{O}^{LL}$  and  $\mathcal{O}^{RL}$  mix with a 3rd operator  $\mathcal{O}^{A(LV)}$   
 $\Rightarrow Z^{AB}$  is a  $3 \times 3$  matrix
- ▶ Exponentially accurate chiral symmetry from Domain Wall Fermions should suppress operator mixing



## NPR

- ▶ Non-perturbative MOM scheme renormalisation of the Rome-Southampton group
- ▶ The renormalised operators are

$$\mathcal{O}_{\text{ren}}^A = Z^{AB} \mathcal{O}_{\text{latt}}^B$$

- ▶ **A** and **B** label the spin structure, eg  $LL$
- ▶  $Z^{AB}$  is the mixing matrix
- ▶  $\mathcal{O}^{LL}$  and  $\mathcal{O}^{RL}$  mix with a 3rd operator  $\mathcal{O}^{A(LV)}$   
 $\Rightarrow Z^{AB}$  is a  $3 \times 3$  matrix
- ▶ Exponentially accurate chiral symmetry from Domain Wall Fermions should suppress operator mixing

## NPR

- ▶ Non-perturbative MOM scheme renormalisation of the Rome-Southampton group
- ▶ The renormalised operators are

$$\mathcal{O}_{\text{ren}}^A = Z^{AB} \mathcal{O}_{\text{latt}}^B$$

- ▶ A and B label the spin structure, eg  $LL$
- ▶  $Z^{AB}$  is the mixing matrix
- ▶  $\mathcal{O}^{LL}$  and  $\mathcal{O}^{RL}$  mix with a 3rd operator  $\mathcal{O}^{A(LV)}$   
 $\Rightarrow Z^{AB}$  is a  $3 \times 3$  matrix
- ▶ Exponentially accurate chiral symmetry from Domain Wall Fermions should suppress operator mixing

## NPR

- ▶ Non-perturbative MOM scheme renormalisation of the Rome-Southampton group
- ▶ The renormalised operators are

$$\mathcal{O}_{\text{ren}}^A = Z^{AB} \mathcal{O}_{\text{latt}}^B$$

- ▶ A and B label the spin structure, eg  $LL$
- ▶  $Z^{AB}$  is the mixing matrix
- ▶  $\mathcal{O}^{LL}$  and  $\mathcal{O}^{RL}$  mix with a 3rd operator  $\mathcal{O}^{A(LV)}$   
 $\Rightarrow Z^{AB}$  is a  $3 \times 3$  matrix
- ▶ Exponentially accurate chiral symmetry from Domain Wall Fermions should suppress operator mixing


## NPR

- ▶ Non-perturbative MOM scheme renormalisation of the Rome-Southampton group
- ▶ The renormalised operators are

$$\mathcal{O}_{\text{ren}}^A = Z^{AB} \mathcal{O}_{\text{latt}}^B$$

- ▶ A and B label the spin structure, eg  $LL$
- ▶  $Z^{AB}$  is the mixing matrix
- ▶  $\mathcal{O}^{LL}$  and  $\mathcal{O}^{RL}$  mix with a 3rd operator  $\mathcal{O}^{A(LV)}$   
 $\Rightarrow Z^{AB}$  is a  $3 \times 3$  matrix
- ▶ Exponentially accurate chiral symmetry from Domain Wall Fermions should suppress operator mixing

We want to calculate the non-perturbative amputated 3-quark vertex function of these operators




$$\mathcal{G}_{abc,\alpha\beta\gamma\delta}^A(p^2) = \epsilon^{abc} (C\Gamma)_{\alpha'\beta'} \Gamma'_{\delta\gamma'} \langle Q_{\alpha'\alpha}^{a'a}(p) Q_{\beta'\beta}^{b'b}(p) Q_{\gamma'\gamma}^{c'c}(p) \rangle$$

where

$$Q_{\alpha'\alpha}^{a'a} = \langle S_{\alpha'\alpha''}^{a'a''}(p) \rangle^{-1} S_{\alpha''\alpha}^{a''a}(p)$$

and  $\Gamma$  and  $\Gamma'$  are the matrices which appear in  $\mathcal{O}^A$

We want to calculate the non-perturbative amputated 3-quark vertex function of these operators




$$\mathcal{G}_{abc,\alpha\beta\gamma\delta}^A(p^2) = \epsilon^{abc} (C\Gamma)_{\alpha'\beta'} \Gamma'_{\delta\gamma'} \langle Q_{\alpha'\alpha}^{a'a}(p) Q_{\beta'\beta}^{b'b}(p) Q_{\gamma'\gamma}^{c'c}(p) \rangle$$

where

$$Q_{\alpha'\alpha}^{a'a} = \langle S_{\alpha'\alpha''}^{a'a''}(p) \rangle^{-1} S_{\alpha''\alpha}^{a''a}(p)$$

and  $\Gamma$  and  $\Gamma'$  are the matrices which appear in  $\mathcal{O}^A$

We want to calculate the non-perturbative amputated 3-quark vertex function of these operators




$$\mathcal{G}_{abc,\alpha\beta\gamma\delta}^A(p^2) = \epsilon^{abc} (C\Gamma)_{\alpha'\beta'} \Gamma'_{\delta\gamma'} \langle Q_{\alpha'\alpha}^{a'a}(p) Q_{\beta'\beta}^{b'b}(p) Q_{\gamma'\gamma}^{c'c}(p) \rangle$$

where

$$Q_{\alpha'\alpha}^{a'a} = \langle S_{\alpha'\alpha''}^{a'a''}(p) \rangle^{-1} S_{\alpha''\alpha}^{a''a}(p)$$

and  $\Gamma$  and  $\Gamma'$  are the matrices which appear in  $\mathcal{O}^A$

We want to calculate the non-perturbative amputated 3-quark vertex function of these operators



$$\mathcal{G}_{abc,\alpha\beta\gamma\delta}^A(p^2) = \epsilon^{abc} (C\Gamma)_{\alpha'\beta'} \Gamma'_{\delta\gamma'} \langle Q_{\alpha'\alpha}^{a'a}(p) Q_{\beta'\beta}^{b'b}(p) Q_{\gamma'\gamma}^{c'c}(p) \rangle$$

where

$$Q_{\alpha'\alpha}^{a'a} = \langle S_{\alpha'\alpha''}^{a'a''}(p) \rangle^{-1} S_{\alpha''\alpha}^{a''a}(p)$$

and  $\Gamma$  and  $\Gamma'$  are the matrices which appear in  $\mathcal{O}^A$



- The renormalization condition in the RI-Mom Scheme is

$$Z_q^{-3/2} Z^{BC} M^{CA} = \delta^{BA}$$

- Where the matrix  $M$  is,

$$M^{AB} = \mathcal{G}_{abc,\alpha\beta\gamma\delta}^A(p^2) P_{abc,\beta\alpha\delta\gamma}^B$$

- and the projection matrices  $P_{abc,\beta\alpha\delta\gamma}^A$  are chosen so that the renormalization condition is satisfied in the free field case where  $Z_q = 1$  and  $Z^{BC} = \delta^{BC}$ .
- $Z^{AB}$  can then be calculated from  $M^{AB}$  using the renormalization condition

- The renormalization condition in the RI-Mom Scheme is

$$Z_q^{-3/2} Z^{BC} M^{CA} = \delta^{BA}$$

- Where the matrix  $M$  is,

$$M^{AB} = \mathcal{G}_{abc,\alpha\beta\gamma\delta}^A(p^2) P_{abc,\beta\alpha\delta\gamma}^B$$

- and the projection matrices  $P_{abc,\beta\alpha\delta\gamma}^A$  are chosen so that the renormalization condition is satisfied in the free field case where  $Z_q = 1$  and  $Z^{BC} = \delta^{BC}$ .
- $Z^{AB}$  can then be calculated from  $M^{AB}$  using the renormalization condition

- The renormalization condition in the RI-Mom Scheme is

$$Z_q^{-3/2} Z^{BC} M^{CA} = \delta^{BA}$$

- Where the matrix  $M$  is,

$$M^{AB} = \mathcal{G}_{abc,\alpha\beta\gamma\delta}^A(p^2) P_{abc,\beta\alpha\delta\gamma}^B$$

- and the projection matrices  $P_{abc,\beta\alpha\delta\gamma}^A$  are chosen so that the renormalization condition is satisfied in the free field case where  $Z_q = 1$  and  $Z^{BC} = \delta^{BC}$ .
- $Z^{AB}$  can then be calculated from  $M^{AB}$  using the renormalization condition

- The renormalization condition in the RI-Mom Scheme is


$$Z_q^{-3/2} Z^{BC} M^{CA} = \delta^{BA}$$

- Where the matrix  $M$  is,

$$M^{AB} = \mathcal{G}_{abc,\alpha\beta\gamma\delta}^A(p^2) P_{abc,\beta\alpha\delta\gamma}^B$$

- and the projection matrices  $P_{abc,\beta\alpha\delta\gamma}^A$  are chosen so that the renormalization condition is satisfied in the free field case where  $Z_q = 1$  and  $Z^{BC} = \delta^{BC}$ .
- $Z^{AB}$  can then be calculated from  $M^{AB}$  using the renormalization condition

$M^{AB}$



- Perform a chiral extrapolation
- Match to the  $\overline{\text{MS}}$  scheme at 2GeV

- This gives

$$U^{\overline{\text{MS}} \leftarrow \text{latt}}(2\text{GeV})_{LL} = 0.662(10)$$

$$U^{\overline{\text{MS}} \leftarrow \text{latt}}(2\text{GeV})_{RL} = 0.664(8)$$

- Additional systematic error of 8% from the matching

- Perform a chiral extrapolation
- Match to the  $\overline{\text{MS}}$  scheme at 2GeV
- This gives

$$U^{\overline{\text{MS}} \leftarrow \text{latt}}(2\text{GeV})_{LL} = 0.662(10)$$


$$U^{\overline{\text{MS}} \leftarrow \text{latt}}(2\text{GeV})_{RL} = 0.664(8)$$

- Additional systematic error of 8% from the matching

- ▶ Perform a chiral extrapolation
- ▶ Match to the  $\overline{\text{MS}}$  scheme at 2GeV
- ▶ This gives
$$U^{\overline{\text{MS}} \leftarrow \text{latt}}(2\text{GeV})_{LL} = 0.662(10)$$
$$U^{\overline{\text{MS}} \leftarrow \text{latt}}(2\text{GeV})_{RL} = 0.664(8)$$
- ▶ Additional systematic error of 8% from the matching



# Summary



- [1] Donoghue 82
- [2] Thomas 83
- [3] Meljanac 82
- [4] Ioffe 81
- [5] Krasnikov 82
- [6] Ioffe 84
- [7] Tomozawa 81
- [8] Brodsky 84
- [9] Hara 86
- [10] Bowler 88
- [11] Gavela 89
- [12] JLQCD 00
- [13] CP-PACS & JLQCD 04
- [14] RBC 07
- [15] RBC 07
- [16] This work

►  $\alpha = -0.0112(12)(22)$


►  $\beta = 0.0120(13)(23)$

## Outlook

- ▶ The direct calculation is currently underway
- ▶ Example: Preliminary results for the  $W_0^{LL}(p \rightarrow \pi^+ + \nu)$ , on the  $16^3 \times 32$  lattice, with valence quark mass  $am_u = 0.03$

## Outlook

- The direct calculation is currently underway
- Example: Preliminary results for the  $W_0^{LL}(p \rightarrow \pi^+ + \nu)$ , on the  $16^3 \times 32$  lattice, with valence quark mass  $am_u = 0.03$



- Writing the proton lifetime as,

$$\Gamma = \text{LEC}^2 \times A_{\text{QCD}} \times A_{\text{GUT}}$$

- Writing the proton lifetime as,

$$\Gamma = \text{LEC}^2 \times A_{\text{QCD}} \times A_{\text{GUT}}$$

- Writing the proton lifetime as,

$$\Gamma = \text{LEC}^2 \times A_{\text{QCD}} \times A_{\text{GUT}}$$

- Writing the proton lifetime as,

$$\Gamma = \text{LEC}^2 \times A_{\text{QCD}} \times A_{\text{GUT}}$$

- Writing the proton lifetime as,

$$\Gamma = \text{LEC}^2 \times A_{\text{QCD}} \times A_{\text{GUT}}$$

- We can put bounds on the GUT scale physics:

Decay Mode	Lifetime bound(yrs)	$A_{\text{GUT}}$ bound ( $M_{\text{GUT}}^{-4}$ )
$p \rightarrow e^+ \pi^0$	$> 8.2 \times 10^{33}$	$< 44$
$p \rightarrow e^+ \pi^0$	$> 8.2 \times 10^{33}$	$< 37$
$p \rightarrow K^+ \bar{\nu}$	$> 2.3 \times 10^{33}$	$< 76$
$n \rightarrow K^0 \bar{\nu}$	$> 1.3 \times 10^{32}$	$< 733$



- For example:  
 $p \rightarrow \pi^0 e^+$  via X Boson Exchange  
Minimal SU(5) SUSY GUT
- $A_{\text{GUT}}$  is given by

$$A_{\text{GUT}} = \frac{g_5^4 A_R^2}{M_X^4} |1 + (1 + V_{ud}^2)^2|$$

- Can put a bound on  $M_X$   
 $M_X > 5 \times 10^{15}$

- For example:  
 $p \rightarrow \pi^0 e^+$  via X Boson Exchange  
Minimal SU(5) SUSY GUT
- $A_{\text{GUT}}$  is given by

$$A_{\text{GUT}} = \frac{g_5^4 A_R^2}{M_X^4} |1 + (1 + V_{ud}^2)^2|$$

- Can put a bound on  $M_X$   
 $M_X > 5 \times 10^{15}$

- For example:  
 $p \rightarrow \pi^0 e^+$  via X Boson Exchange  
Minimal SU(5) SUSY GUT
- $A_{\text{GUT}}$  is given by

$$A_{\text{GUT}} = \frac{g_5^4 A_R^2}{M_X^4} |1 + (1 + V_{ud}^2)^2|$$

- Can put a bound on  $M_X$   
 $M_X > 5 \times 10^{15}$